

The phone rang and I heard an unfamiliar voice. She spoke of **"the most horrible thing she had ever seen!"**

She was a Registered Nurse working in an Emergency Room in a near-by hospital. She had witnessed not one, but two injured women brought in with near death injuries.

The cases were similar in that they both had late term abortions with their babies at 25 weeks.

In both patients, their wombs were torn and the babies had been forced into their abdominal cavities.

The decimation of the insides of these two women was caused by violent contractions brought on by a medication manufactured by

Pfizer called MISOPROSTOL also known as CYTOTEC.

The extent of the violence was enormous!

The first woman had to have an emergency colostomy to save her life. She was in a coma for 4 days and in the hospital for 9 days. The second woman had to have a hysterectomy to save her life and was in the hospital for 9 days. From 6 to 36 weeks, almost all women having an abortion are administered **MISOPROSTOL/CYTOTEC.**



We Care.

**Visit our website
to keep updated
on how we can help you.**



WAKEUP

www.wakeup.mom



Abortion occurs with

MIFEPREX
(RU-486)

...BLOCKS PROGESTERONE

&

MISOPROSTOL
(CYTOTEC)

...INDUCES CONTRACTIONS

FATAL

Details Enclosed

MISOPROSTOL OR CYTOTEC IS A FDA APPROVED MEDICATION USED TO TREAT ULCERS.

It was found as a side effect to have caused pregnant women to miscarry.

THE USE OF MISOPROSTOL OR CYTOTEC IS **NOT** FDA APPROVED FOR USE DURING ABORTIONS.

It is "off label" usage and can cause serious, even fatal side effects.

MISOPROSTOL OR CYTOTEC

- Causes violent and uncontrollable uterine contractions in the clinic or at home.
- Is used in most abortion procedures.
- Is used in conjunction with the RU-486, also known as the medical abortion or pill abortion. In second and third trimester abortions, it is also given to soften the cervix.
- This medication may have been given to you to take home or administered at the clinic.
- This medication can be taken orally, buccally (dissolved between cheek and gum), inserted vaginally or rectally.

THE USE OF MISOPROSTOL OR CYTOTEC CAN CAUSE:

- Intense Pain
- Severe Cramping
- Severe Contractions
- Nausea
- Vomiting
- Diarrhea
- Fever and Chills
- Infection
- Fainting
- Emergency Intervention
- Uterine Lacerations
- Rupture of Uterus
- Blood Transfusion
- Death

Women all over the United States are suffering complications from the use of

MISOPROSTOL OR CYTOTEC.

These complications and injuries have catastrophic life changing outcomes.

*For Help. . .
Simply Call Us.*

- If you have been injured during an abortion and are not sure if this medication was the cause, call us for **HELP**.
- If you know this medication caused you a complication and are interested in joining a class action lawsuit against its manufacturer.